

# Dusty Air Contributing to Disease in Africa

By

David J. Craig

|

Spring 2014



Dust storms like this one in Sudan appear to be making people vulnerable to meningitis. Photo: Reuters / Mohamed Nureldin Abdallah

Among the hardships wrought by the dry season in sub-Saharan Africa, which lasts from November to May, are outbreaks of bacterial meningitis. In a bad year, this disease, which attacks the thin lining of the brain and spinal cord, can affect tens of

thousands of people in a semiarid section of Africa that covers large parts of Sudan, Chad, Niger, Nigeria, and Mali.

Until now, scientists weren't sure why the disease is seasonal. Columbia climate scientists led by Carlos Pérez García-Pando say they've found an answer. By studying twenty years' worth of climate data and health records in Niger, they have concluded that a good predictor of meningitis rates is the amount of dust kicked up by winds.

The scientists suspect that people who breathe in lots of dust are vulnerable to meningitis because they have tiny abrasions in their throats. They say that African officials ought to adjust where they distribute vaccines each year, based partly on wind conditions, so that they can respond faster to outbreaks.

Read more from [David J. Craig](#)



[Guide to school abbreviations](#)

# TAKE THE COLUMBIA ALL-ALUMNI SURVEY

Complete the survey  
by June 5.

50 randomly selected  
survey participants  
will receive a  
Columbia sweatshirt!

Shape the alumni  
experience.

[alumni.columbia.edu/survey2026](https://alumni.columbia.edu/survey2026)

 COLUMBIA UNIVERSITY  
IN THE CITY OF NEW YORK

[All categories >](#)

[Read more from \*\*David J. Craig\*\*](#)